
## **CPWR**

## **OPIOIDS AND MENTAL HEALTH**

17 MAY 2023


Christina Trahan Cain, CIH
CPWR Executive Director
NABTU Safety and Health Director





Industry sectors with opioid-related overdose death rates significantly higher than the average rate for all workers, Massachusetts workers, 2011-2015, n=4,302

## Suicide rates among male U.S. workers, by major occupation group, 2015



# What has the construction industry done about it?

## NABTU Opioid Task Force

NABTU President Sean McGarvey established

- 14 international union reps
- Employers and employer reps
- BTCs, Insurers, and Government partners

Adopted a public health model to address the problem





## Public Health Model

Tertiary Prevention

SUD Treatment and Recovery

Secondary Prevention Treatment Alternatives to Opioids

**Primary Prevention** 

**Prevent Pain** 

**Prevent Injuries** 



## NABTU 2020 Resolution

"Support for Efforts to Reduce Pain. Opioid Use, **Opioid Overdose** and the Number of Deaths by Suicide in the Construction Industry"

## Resolution No. 4

Re: Support for Efforts to Reduce Pain, Opioid Use, Opioid Overdose and the Number of Deaths by Suicide in the Construction Industry

Submitted by: Governing Board of Presidents

WHEREAS, in the United States, the combined number of deaths among Americans from suicide and unintentional overdose increased from 41,364 in 2000 to 110,749 in 2017. Among occupations, the construction industry has the second highest rate of both death by suicide and opioid overdoses; and

WHEREAS, among all workers, unintentional overdoses have increased 420% between 2011 and 2018 and have increased 930% for construction workers during the same time period; and

WHEREAS, the injury rate for construction workers is 77 percent higher than the national average for other occupations; and

WHEREAS, injured construction workers are not likely to be offered modified duty, more likely to suffer pain and job loss and/or pressure to return to work quickly, and experience financial stress, all of which contribute to depression and increased opioid usage; and

WHEREAS, studies in Ohio and Massachusetts, both high union density states, found that construction workers had a disproportionate number of opioid overdose deaths; and

WHEREAS, chronic pain, depression, and opioid use are associated with increased risk of suicide and construction ranks as the industry with the second greatest number of suicides; and

WHEREAS, injured workers receiving workers' compensation have been more likely to receive opioid prescriptions for general pain and to recover from medical procedures than those with non-work-related injuries undergoing the same procedure;

THEREFORE, BE IT RESOLVED, that North America's Building Trades Unions and all Building Trades Councils fully endorse combatting opioid-related deaths and deaths by suicide in the construction industry by taking these measures to prevent pain, educate the industry, and provide support to members:

· Promote programs and ergonomic equipment on job sites that reduce musculoskeletal disorders and traumatic injuries-work shouldn't hurt.

· Work to destigmatize substance use and mental health disorders through culturally and linguistically appropriate services, education and awareness, with members, leadership,

 Educate members about the problems and limitations of opioids for long term treatment of injuries and chronic pain resulting from construction work—and informing them about non-opioid alternatives to pain management treatment.

 Mandate all apprentice and/or trainee members to complete a training program designed to increase awareness of work-related injuries associated with opioid use.

- Design International and Local Taft-Hartley health funds to provide members with best in class benefits to promote behavioral health and substance use disorder benefits and alternative treatment for pain. For example, evaluate benefit designs to ensure compliance with the Mental Health Parity Act and promote medication-assisted treatment (MAT), including opioid treatment programs (OTPs), that are combined with behavioral therapy and medications to treat substance use disorders. Consider covering services that are non-traditional but effective for pain management such as acupuncture, massage, and physical therapy. Ensure that pharmacy benefit managers are offering clinical management programs such as step-therapy, quantity level limits, and clinical prior-authorization to ensure that best practices are followed.
- Support Nalaxone trainings for members.
- · Develop peer educator programs to connect affected workers with substance use disorder treatment and mental health support.
- · Publicize available behavioral health resources, inclusive of member/employee assistance programs, peer programs, and counseling and treatment resources, available through building trades unions and health and welfare funds.
- Support members at all steps in their path to recovery from substance use disorder or behavioral health issues. This includes pre-treatment, treatment and long-term
- · Educate members and provide resources on suicide prevention and awareness.
- · Encourage organizations to help develop and support workplace policies and programs that promote rehabilitation and return-to-work opportunities.




## **CPWR** Resources



## Projects to Prevent Opioid Use

- Communications Report on Primary Prevention
- Training Programs
- Peer Advocacy Report
- Data Reports
- Physicians Alert
- Aids to Reduce Stigma



Source: <a href="https://www.cpwr.com/wp-content/uploads/KF2021-opioid-related-harms-prevention.pdf">https://www.cpwr.com/wp-content/uploads/KF2021-opioid-related-harms-prevention.pdf</a>



## FrameWorks Communication Solutions

- Link causes and consequences to build support for structural solutions.
- Use the *Upstream/Downstream* metaphor to explain prevention.
- Appeal to the Value of Investment in messages to construction industry.
- Choose concrete examples to illustrate what effective interventions look like.
- Provide the context needed to interpret unfamiliar concepts and data.
- Explicitly name who or what is responsible for the problem or taking action to fix it.
- Emphasize systemic solutions to expand thinking beyond individual-level interventions.



## Opioid Awareness Training

- Created an opioid hazard awareness training on behalf of North America's Building Trades Unions
  - Improve knowledge about opioids and related substance use and mental health
  - Inspire and motivate trainees to act
- Piloted and Evaluated
- Shortened and digitally-optimized training in 2020, updated 2022
- Free for download: <u>www.cpwr.com</u>



## Suicide Training/Mental Health Program

## 9 discussion-based activities to be used in training curricula:

- 1. Understanding Member Assistance Programs
- 2. NABTU Resolution: Support for Efforts to Reduce Pain, Opioid Use, Opioid Overdose and the Number of Deaths by Suicide in the Construction Industry
- 3. Motivational Interviewing
- 4. Understanding Upstream Prevention
- Basic Suicide Prevention
- 6. Health Maintenance
- 7. Drug Testing Debate
- 8. A Suicide on Site: Getting the Story Right
- 9. Roach Coach Health Conversation



## Peer Advocacy in the Construction Industry

- Interviewed Key Informants from the NABTU Opioid Task Force
  - Union Response to Opioid Crisis
- 7 of 13 Interviewees Discussed Peer Advocacy
  - Themes about Peer Advocacy Included:
    - Barriers Stigma, Buy-In, Trust
    - Planning
    - Design
    - Recovery

PEER ADVOCACY FOR CONSTRUCTION WORKERS STRUGGLING WITH SUBSTANCE USE AND MENTAL HEALTH







## **Data Center Reports**

Overdoses

Opioid Use

Mental Health

during COVID



## Overdose Fatalities at Worksites and Opioid Use in the Construction Industry

Xiuwen Sue Dong, DrPH\*, Raina D. Brooks, MPH, Chris Trahan Cain, ClH

## Foreword

Construction workers are among the segments of the U.S. population opioids have hit hardest. Recent state-level studies of opioid overdose deaths show that construction workers are six to seven times more likely to die of an overdose than workers in other professions. The impact of opioids to our field led us to make it the focus of this Quarterly Data Report.

Section 1 examines a small subset of construction workers who died of an overdose: those who died on a worksite. These are figures for which we have national data, but there is not equivalent national data yet about how many of the 130 Americans who die each day from an opioid overdose work in construction.

This report also reveals other gaps in our understanding of the impact of opioids on construction workers. For example, Section 2 contains the surprising finding that the percentage of construction workers who used prescribed opioids, on average, is slightly lower than workers in all industries combined. Our assumption before conducting this analysis was the reverse, given that construction has one of the highest injury rates of all industries, particularly musculoskeletal disorders that often result in chronic pain and long-term pain management. One possible explanation for this counter-intuitive finding: construction workers are less likely to have health insurance than workers in other major industry sectors, and so they may be less likely to receive a prescription for opioids than workers in other sectors.

While the impact of opioids on the construction industry and its workers is becoming clearer, there remains much we need to learn to understand and respond to the damage they are causing. We look forward to receiving your feedback on this important report and working collectively to minimize the impact opioids are having on workers, their families, the industry, and society overall.

Chris Trahan Cain Executive Director CPWR.

## KEY FINDINGS

- Unintentional overdose fatalities in the construction industry jumped from 7 deaths in 2011 to 65 deaths in 2018, a nine-fold increase in eight years.
- Between 2011 and 2017, one in four (25.3%) construction workers with work-related injuries used prescribed opioid pain relievers. compared to approximately one in ten (8.9%) of their counterparts who were not injured.
- Older construction workers were more likely to use prescribed opioid pain relievers, while younger. construction workers were more likely to use illicit
- Uninsured construction workers were less likely to use prescribed opioid pain relievers, but more likely to use illicit drugs than their insured counterparts.









## Construction Worker Mental Health During the COVID-19 Pandemic

Samantha Brown, MPH, Amber Brooke Trueblood, DrPH, William Harris, MS, Xiuwen Sue Dong, DrPH1

## **OVERVIEW**

Anxiety and depression symptoms significantly worsened nationwide during the COVID-19 pandemic. Construction workers already suffer from an increasing and alarmingly high suicide rate, making it particularly important to understand mental health in the industry during the pandemic. To support that goal, this Data Bulletin examines self-reported symptoms of anxiety and depression in the population using the National Health Interview Survey (NHIS) from 2011 to 2018 and in 2020,2 focusing on patterns and changes during the pandemic. Anxiety and depression were measured for construction workers by A) feelings of anxiety or depression at least once a month; and B) feelings of anxiety or depression at least once a week, or associated medication use. (see the Definitions section at the end of the report for detailed criteria). Differences in the frequency or level of anxiety/depression between 2019 and 2020 were measured in a subsample of construction workers who were interviewed in both years. Anxiety/ depression was compared across3 worker demographics, socioeconomic status, and health indicators (i.e., health status, alcohol use, opioid use, and health insurance coverage). Due to the survey methodology changes in 2020 and fewer respondents during the pandemic, the sample size of some subgroups is relatively small.4



<sup>1</sup>Correspondence to:datacenter@epwr.com.

<sup>2</sup>No industry and occupation information in the 2019 survey due to the questionnaire redesign.

Statistical significance is not discussed in the text but is provided in the associated charts.

Frequencies of anxiety/depression are small (n < 30) for some subgroups in certain charts (see chart footnotes). Readers are advised to use related results with caution. Numbers in text and charts were calculated by the CPWR Data Center.

## THIS ISSUE

This issue examines anxiety and depression symptoms or medication use among construction workers before and during the COVID-19 pandemic. comparing differences by demographics, socioeconomic status, and health indicators.

## KEY FINDINGS

Construction workers feeling anxious at least once per month rose 20% between 2011 and 2018.

In 2020, the prevalence of anxiety/depression (based on feelings or medication) in workers was 15%, and was particularly high in those who were age 18-34 (18%), female (24%), living below the poverty line (18%), or working part-time (19%).

Charts 4-6

In 2020, symptoms or medication use for anxiety/ depression were almost three times higher in workers who used prescription opioids in the past year compared to those who did not (39% versus 14%).

Among workers who were surveyed in both 2019 and 2020, 43% had increases in the frequency or level of anxious/ depressed feelings between vears, with increases more common in those who were age 18-54 (46%), female (50%), or had a family income below the poverty line (61%).

Charts 8-10

## NEXT DATA BULLETIN

EmploymentTrends and Projections in Construction



Source: https://www.cpwr.com/research/data-center/data-reports/

## **CPWR Physicians'/Providers' Alerts Document**

## **LEVEL 2 Prevention: Avoid Opioids**

Avoid

Avoid long-term opioid prescriptions

**Avoid** 

Avoid combined prescriptions (tranquilizers + muscle relaxants + painkillers)

Advocate

Advocate for good care, including nonopioid treatment

## Physicians'/Providers' Alert:

## Pain Management for Construction Workers

This Alert was developed to help ensure that all construction workers who visit a doctor or other healthcare provider because of pain from an injury are aware of treatment options and understand the potential risks of addiction associated with wine practication options. Blacks:

- (1) read and print this Alert;
- (2) keep the "Tips for Talking with Your Doctor"; and
- (3) fill in the "To My Doctor" form and give it to your doctor to include in your medical records

## Tips for Talking with Your Doctor: What You Need to Know Before Accepting an Opioid Prescription

Opicids, such as fentany (Duragesic\*), hydrocodone (Vicodin\*), oxycodone (OxyContin\*), oxymorphone (Opana\*), hydromorphone (Dilaudid\*), meperidine (Demerol\*), diphenoxylate (Lomotil\*), tramadol, buprenorphine (e.g., Suboxone\*), morphine, and codeine are often prescribed to help manage pain. In addition, new drugs are entering the market place, such as Dsuvia\*, which are considered even more addictive. Since these medications can be addictive, they should only be used if other treatment options are not effective. When prescribed, they should be used for the shortest time possible, be closely monitored, and include courseling.

Talk to your doctor about treatment options and how the medication may affect you. Remember to tell your doctor

- If you have been or are being treated for another health issue or have been prescribed other medications by another doctor.
- If you have a history of addiction to tobacco, alcohol or drugs, or if there is a history of addiction in your family.
  About your work environment. Let your doctor know that 1) taking opioids on the job can be a safety hazard because they can make you drowsy, and 2) testing positive for some drugs, even when prescribed for pain, can negatively impact employment opportunities. Some employers have expanded panels of drugs they test employees for, which are regularly reviewed and udadated. The Department of Transportation's drug test posel, for exampled.
  - Opioids (codeine, morphine, 6-AM (heroin), hydrocodone, hydromorphone, oxycodone, oxymorphone)
  - Phencyclidine
- Marijuana (THC)
- Cocaine
- Amphetamines (amphetamine, methamphetamine, MDMA, MDA)

Before accepting a prescription for one of the medications listed earlier or another opioid, ask your doctor/healthcare

- 1. Can my condition be effectively treated without opioid medication? If yes, what would the treatment involve?
- [If prescribed an opioid and are taking other medications] Will the opioid medication interfere with other medications that I'm currently taking?
- Are there potential side effects from the opioid medication prescribed? If yes, how can I reduce the risk of side

## emember:

NEVER share medications or store medications where others will have access.

ALWAYS safely dispose of medications. Look for a medicine disposal center near you (often at your local pharmacy)

### To learn more visi

- CPWR Opioid Resources website <a href="https://www.cpwr.com/research/opioid-resources">https://www.cpwr.com/research/opioid-resources</a>
- Substance Abuse and Mental Health Services Administration (SAMHSA) <a href="https://www.samhsa.gov/">https://www.samhsa.gov/</a> or call their confidential national hotline 1-800-662-HELP (4357)
- Facing Addiction's online Addiction Resource Hub <a href="https://resources.facingaddiction.org/">https://resources.facingaddiction.org/</a>

<sup>3</sup> Source: U.S. Department of Transportation. (2018). DOT 5 panel notice. https://www.transportation.gov/odapc/DOT\_5\_Panel\_Notice\_2011

CPWR THE CENTER FOR CONSTRUCTION

Source: <a href="https://www.cpwr.com/wp-content/uploads/publications/publications\_handouts-and-toolbox-talks\_physicians-alerts\_opioids.pdf">https://www.cpwr.com/wp-content/uploads/publications/publications\_handouts-and-toolbox-talks\_physicians-alerts\_opioids.pdf</a>



## Jobsite Opioid Resources



## **OPIOID DEATHS** IN CONSTRUCTION



## Why Are Construction Workers at Risk?

- The construction industry has one of the highest injury rates compared to other industries.
- > Opioids are often prescribed to treat the pain caused by these
- Long-term opioid use can make people more sensitive to pain and decrease the opioid's pain-reducing effects.





According to the CDC, 1 out of 4 people prescribed opioids for long-term pain become addicted.





## **Injured Construction** Workers Often...

- Cannot continue to work while injured. Suffer a loss in income. Even if an injured worker receives workers' compensation, it is often not enough to make up for
- Experience anxiety, stress, and depression, which can add to



## **Overdose Deaths Are** On the Rise.

- ▶ In 2020 alone, there were more than 93,000 overdose deaths in the U.S.—nearly 75% of which involved an opioid. 1 Opioid-related overdose deaths increased 36.7%
- These increases were exacerbated by the COMD-19 pandemic because of isolation, loss of loved ones, stress, unemployment, precarious housing, decreased access to treatment services, and more 3
- Construction workers are significantly more likely to die from an opicid overdose than the average worker - studies in both Chio and Massachusetts, for example, showed they were seven
- Additionally, a past study has shown that more than half of those who died from an overdose had suffered at least one

## **Protect Yourself!**



Work shouldn't hurt. Your employer should be committed to a safe job site, and you should use safe practices. Together these reduce the risk of injuries and therefore the need for pain



Talk to a Doctor Opioids are addictive and can have side



- Other forms of pain medication that are not addictive and have fewer side effects
- Other forms of pain management such as physical therapy or acupuncture Opicids should be the last option to treat your pain. If opicids are prescribed they should be used for the shortest possible time. Safely dispose of any unused medications.



## Get Help

Opioids change how your brain works, triggering one part of it to demand more opioids and changing another part of it so it's harder to resist." Check with your union or employer to find out if they have a program to help, such as:

- An employee assistance program (EAP); or
- Member assistance program (MAP).

If you're having trouble stopping using opioids, check with your union or your doctor for help to find the best addiction treatment option for you.

nber addiction is an illness that can be treated Call this confidential national hotline: 1-800-662-HELP (4357)

### If you or someone you know needs help:

- Contact the Substance Abuse and Mental Health Services Administration at https://www.samhsa.gov.or.cal their confidentia national holline: 1-800-662-HELP (4357).
- Visit the Recovery Resource online hub:
- Contact your union.
- Find a list of common opioids at:
- Give your doctor the Physicians' Alert on Pain Management among Construction Workers from

To receive copies of this Hazard Alert and cards on other topics: call 301-578-8500 email cpwr-r2p@cpwr.com

Find out more about





## **Opioid Deaths** in Construction

Construction work can result in painful injuries that are sometimes treated with prescription opioids. One in four people prescribed opioids for long-term pain become addicted and opioid-related deaths

## Chris' Story

Chris strained his back after lifting heavy materials. He tried to ignore the pain, but it wouldn't go away. Chris went to the doctor and was prescribed an opioid to treat the pain. The pills reduced the pain, but his back never got better. Chris found that he needed the pills to make it through the day. Eventually, his doctor refused to give him another prescription. Chris went to another doctor and got a new prescription. Over time his job performance and family life began to suffer. Chris went back to his doctor and asked for help. His doctor helped him to find treatment for his opioid addiction. Chris is now in recovery and using a non-addictive treatment for

- Have you known someone addicted to opioids?
- or If a worker is injured and in pain, what should he or she do to avoid becoming addicted to

## Remember This

- Your employer must provide a safe work environment to prevent injuries. If you see a hazard on the job, report it to your supervisor or
- Follow safe work practices to prevent injuries, such as getting help when lifting heavy
- If you are injured, talk to your doctor about non-addictive medications or physical therapy to treat the pain.
- Opioids should be the last option, and if prescribed used for the shortest time possible.
- Addiction is an illness that can be treated. Get help if you find you are dependent on pain medication to get through the day.
- Check with your union or employer to find out if they have a program to help, such as an employee assistance program (EAP) or member assistance program (MAP).
- Call this confidential national hotline to find out about treatment options near you 1-800-662-HELP (4357) or go online at https://resources.facingaddiction.org.

\*Centers for Disease Control & Prevention, Promoting Safer and More Effective Pain Management. https://www.cdc.gov/drugoverdose/pdf/Guidelines\_Factsheet-Patients-a.pdf



Source: https://www.cpwr.com/research/research-to-practice-r2p/r2plibrary/other-resources-for-stakeholders/mental-healthaddiction/opioid-resources/ 16



## Infographic



Construction work can result in painful injuries that are often treated with prescription opioids.

Opioids are addictive and should be the last option to treat your pain. Talk to your doctor about non-addictive medications.





**REMEMBER:** Addiction is an illness that can be treated.


Call this confidential national hotline:

1-800-662-HELP (4357)

Visit: Facing Addiction — https://resources.facingaddiction.org/







CPWR THE CENTER FOR CONSTRUCTION RESEARCH AND TRAINING

Source: https://www.cpwr.com/research/research-to-practice-r2p/r2plibrary/other-resources-for-stakeholders/mental-health-addiction/opioidresources/dont-be-a-statistic-protect-yourself-from-an-opioid7overdose/



## Jobsite Suicide Prevention Resources



## Suicide Prevention



## The Data

Suicide rates in the U.S. have increased in recent years, and it has been the 10<sup>th</sup> leading cause of death since 2008. In 2018<sup>th</sup> alone, there were:

## 48,344 deaths – an average of 132 per day or 1 every 11 minutes.

Suicide can affect anyone. According to the Centres for Disease Control and Prevention (CDC), construction has one of the highest suicide rates compared to other industries: "There is no simple areaser to why this increase has occurred, particularly among construction workers However, there are steps that workers and employers can take to recognize the warning signs and help revenent sicilities."

## **Recognize the Warning Signs**

According to mental health professionals, the following are common warning signs that a person may be thinking about suicide1:

Talking about wanting to die, guilt or shame, or being a burden to others.

### Feeling:

 Empty, hopeless, or having no reason to live; extremely sad, anxious, agitated, or angry; unbearable emotional or physical pain.

### Behavior:

- Planning or researching ways to die; buying a gun; withdrawing from friends, family, or activities, saying goodbye, giving away possessions, or making a will.
- Agitation or rage increased conflict among co-workers<sup>2</sup>; extreme mood swings; changes in personality or neglecting their appearance.
- Taking dangerous risks, such as increased alcohol or drug use or driving recklessly; eating or sleeping more or less; increased tardiness and absenteeism from work?.

Sources: 1) National Institute of Mental Health, Warning Signs of Suicide. https://bib/p2XXXZ62 2) Construction Francial Management Association. Face Suicide Vierning Signs Before It's Too Late. https://bib. by/34/55M



Find out more about construction hazards.

To receive copies of this Hazard Alert and cards on other topics call 301-578-8500 or

call 301-578-8500 or email cpwr-r2p@cpwr.com 8484 Qeorgia A Suts 1000

CPWR Subs 1000
Sher Spring, MD 3
Sher Spring, MD

## Help Prevent Suicide...

## 1 Reach Out

If you notice the warning signs of suicide in someone you know, talk to them – start a conversation. Ask them about a specific warning sign you've noticed. For exemple, "I've noticed lately that you are sitting alone at lunch and avoiding all of us while we re at work, and are concerned." You may feel uncominative, but the best way to find out if someone is having suicide thought is to sak them directly. "Are you thinking about suicide?" Asking this will not put the idea into their head or make it more likely that they will attempt suicide.

If the answer is "Yes," do not leave them alone and get help.

Source: California Mental Health Sovices Authority (CalMHSA). Know the Signs https://www.suiciteispreventable.org

## 2 Respond

When taking to someone who may be thinking about suicide, take what they say seriously. Listen without judgement, and express concern and support. Be direct. Talk openly and matter-of-facity about suicide. Do not ask questions encouraging them to deny their feelings such as: "Bour en of thinking about suicide, are you?" Reassure them that he'd is available.

### DO NOT:

Tell the person to do it; debate the value of living or argue that suicide is right or wrong; minimize their problems by saying things like "You'll get over it," Toughen up," or "You're fine"; promise to keep their thoughts about suicide a secret.

Source: National Suicide Prevention Lifetine. Help Someone Elea. https://bit.ly/2xapkPt.

## 3 Connect

Encourage the person to see a mental health professional. Call the National Suicide Prevention Lifeline for advice and referrals, or help them locate a treatment facility or program.

Stay in touch with them after a crisis to see how they are doing. Remind

## ▶ YOU ARE NOT ALONE. THERE IS HOPE. SUICIDE IS NOT THE ANSWER.

If someone is in immediate danger, call 911, take them to a nearby emergency room, call the **National Suicide Prevention Lifetine** at 1-800-273-8255, or reach out to the Crisis Text Line by texting "HELLO" to 741741 to connect with a crisis counselor.

### To Learn More About Preventing Suicides, Visit:

CPWR - Suicide Prevention
Resources:
https://finyurl.com/Suicide-Prevention

https://tinyurl.com/Suicide-Prevention

Construction Industry Alliance for
Suicide Prevention:

American Foundation for Suicide Prevention: https://afsp.org/

### If You or Someone You Know Needs Immediate Help, Contact: The National Suicide Prevention

## Lifeline: Provides free and confidential support from

Provides free and confidential support fro trained counselors 24/7.

 Call 1-800-273-TALK (8255)
 Use the online Lifeline Chat at: https://bit.lv/2WG0bnl

Crisis Text Line: Text "HELLO" to 741741 for free, 24/7 confidential support.



## Suicide Prevention in Construction



Over the last several years, the rate of suicide has increased, and it is now the 10th leading cause of death in the U.S. The construction industry has one of the highest suicide rates compared to other industries. Although there is no simple reason for this increase, learning the warning signs and how to reach out for help could save your life or the life of a co-worker.

## John and Matt's Story

Matt noticed that his co-worker John was acting differently, becoming easily upset, not following safe practices on the job, and eating lunch alone. He recognized that these may be the warning signs of suicide. During lunch he called a crisis hotline and asked for advice on how to talk to John and get him help. At the end of the day, Matt approached John and said that he had noticed a change in his mood and behavior lately and was concerned. He asked John if he has had any thoughts of suicide and told him about the hotline. John was angry at first, but then admitted that he had been feeling depressed and is relieved that he can talk to someone about it. Matt convinced John to call the crisis hotline. John is now getting the help he needs, and Matt is confining to provide support.

- thoughts or died by suicide?
- What are examples of the warning signs of suicide?
- How can we help a co-worker when there are warning signs of suicide?

## Remember This

Recognize the Warning Signs:

## Talking about:

- Wanting to die
- Guilt or shame Being a burden to others

### Feeling:

- Empty, hopeless, trapped, or having no reason to live
- Extremely sad, anxious, agitated, or angry
- Unbearable emotional or physical pain

## Behavior:

- Planning or researching ways to die; purchasing a gun
   Withdrawing from friends, family, or activities, saying goodbye, giving away possessions, or making a will
- Agitation or rage increased conflict among co-workers
- Extreme mood swings
   Changes in personality or neglecting their appearance
- Taking dangerous risks, such as increased alcohol or drug use or driving recklessly
- Eating or sleeping more or less
- Increased tardiness and absenteeism from work
- If someone you know is showing any of these signs, don't ignore them. Start a conversation. The best way to find out if a person is having suicidal thoughts is to ask directly.
- Listen without judgement and express concern and support.
   Reassure them that help is available.
- DO NOT tell someone to do it, debate the value of living, or argue that suicide is right or wrong.
- > NEVER promise to keep their thoughts about suicide a secret.
- Encourage the person to see a mental health professional or help them locate a treatment facility.
- If you believe someone is in immediate danger, call 911, take them to a nearby emergency room, call the National Suicide Prevention Lifeline at 1-800-273-8256, or reach out to the Crisis Text Line by texting "HELLO" to 741741 to connect with a crisis
- Stay in touch with them after a crisis to see how they are doing.

If you or someone you know needs immediate help, contact the National Suicide Prevention Lifeline at 1-800-273-TALK (8255), use the online Lifeline Chat, or text "HELLO" to 741741 to connect with a crisis counselor. They provide free and confidential support with trained counselors 24/7.

_	ow can we stay safe today? hat can we do today to help prevent suicide?
) 1.	
2.	
(	

©2020, CPWR.The Center for Construction Research and Training. All rights reserved. CPWR is the research and training arm of NABTU. Production of this document was supported by cooperative agreement OH 009762 from the National institute for Occupational Safety and Health (NIGSN). The contents are solely the responsibility of the authors and do not necessarily represent the official viscous for NICSH.



Source: https://www.cpwr.com/research/research-to-practice-r2p/r2p-

<u>library/other-resources-for-stakeholders/mental-health-addiction/suicide-</u>

prevention-resources/





## Infographic



If you or someone you know needs immediate help, contact the National Suicide Prevention Lifeline at 1-800-273-TALK (8255) or text "HELLO" to 741741 to connect with a crisis counselor.

62020, CPWR-The Center for Construction Research and Training, All rights reserved, CPWR is the research and training arm of NABTU. Production of this document was supported by cooperative agreement OH 000702 from the National Institute for Occupational Safety and Health (NIOSH). The contents are solely the responsibility of the authors and do not necessarily represent the official views of NIOSH.





If you or someone you know needs immediate help, contact the National Suicide Prevention Lifeline at 1-800-273-TALK (8255) or text "HELLO" to 741741 to connect with a crisis counselor.

@2020, CPVM-The Center for Construction Research and Training, All rights reserved. CPVM is the research and training arm of NABTU. Production of this document was supported by cooperative agreement 0H 009782 from the National Institute for Occupational Safety and Health (NIOSH). The contents are solely the responsibility of the authors and do not necessarily represent the official views of NIOSH.



Source: <a href="https://www.cpwr.com/research/research-to-practice-r2p/r2p-library/other-resources-for-stakeholders/mental-health-addiction/suicide-prevention-resources/suicide-prevention-infographics/">https://www.cpwr.com/research/research-to-practice-r2p/r2p-library/other-resources-for-stakeholders/mental-health-addiction/suicide-prevention-resources/suicide-prevention-infographics/</a>



## Ongoing Work



## **Bullying Survey**

- Apprentices
- In partnership with:
  - International Union,
  - National Employer Association, and
  - National Training Fund



## CPWR August 2022 Opioids/Suicide Workshop





## Workshop Outcome

## Four Ongoing Workgroups

- Training and Education
- Changing the Culture and Stigma Reduction
- Injury Prevention and Workplace Stress
- Peer Support

CPWR will support the workgroups and coordinate with the NABTU Opioid Task Force moving forward.



## CPWR.com

- All <u>CPWR</u> free resources and more
  - SAMHSA Treatment Locator
  - National Suicide & Crisis Lifeline
  - CIASP Website Links
  - NIOSH
  - CDC



